

The loopy UV line profiles of RU Lupi: accretion, outflows, and fluorescence

Gregory J. Herczeg¹, Frederick M. Walter², Jeffrey L. Linsky¹, Gösta F. Gahm³, David R. Ardila⁴, Alexander Brown⁵, Christopher M. Johns-Krull⁶, Michal Simon², Jeff A. Valenti⁷

ABSTRACT

We present far-ultraviolet spectra of the classical T Tauri star RU Lupi covering the 912–1710 Å spectral range, as observed by the *HST*/STIS and *FUSE* satellites. We use these spectra, which are rich in emission and absorption lines, to probe both the accreting and outflowing gas. Absorption in the Ly α profile constrains the extinction to $A_V \sim 0.07$ mag, which we confirm with other diagnostics. We estimate a mass accretion rate of $(5 \pm 2) \times 10^{-8} M_\odot \text{ yr}^{-1}$ using the optical-NUV accretion continuum. The accreting gas is also detected in bright, broad lines of C IV, Si IV, and N V, which all show complex structures across the line profile. Many other emission lines, including those of H₂ and Fe II, are pumped by Ly α . RU Lupi's spectrum varies significantly in the FUV; our STIS observations occurred when RU Lupi was brighter than several other observations in the FUV, possibly due to a high mass accretion rate.

Subject headings: accretion, accretion disks — circumstellar matter — line: identification — stars: individual (RU Lupi) — stars: pre-main sequence — ultraviolet: stars

1. INTRODUCTION

Classical T Tauri stars (CTTSs) are likely prototypes of the young Sun when it was accreting material from its circumstellar disk. In the standard magnetospheric accretion model, the magnetic

¹JILA, University of Colorado and NIST, Boulder, CO 80309-0440; gregoryh@casa.colorado.edu, jlin-sky@jila.colorado.edu

²Department of Physics and Astronomy, Stony Brook University, Stony Brook NY 11794-3800; fwalter@astro.sunysb.edu, msimon@astro.sunysb.edu

³Stockholm Observatory, AlbaNova, SE - 106 91 Stockholm, Sweden; gahm@astro.su.se

⁴Department of Physics & Astronomy, Johns Hopkins University, Baltimore MD 21218; ardila@pha.jhu.edu

⁵Center for Astrophysics and Space Astronomy, University of Colorado, Boulder CO 80309-0389; ab@casa.colorado.edu

⁶Department of Physics & Astronomy, Rice University, Houston TX 77005-1892; cmj@rice.edu

⁷Space Telescope Science Institute, Baltimore MD 21218; valenti@stsci.edu

field truncates the disk at a few stellar radii. Material from the disk accretes onto the star along the magnetic field lines and is shocked as it decelerates at the stellar surface. A fast outflow is likely associated with the accretion process, and a slow outflow probably emanates from the disk (see discussion in, e.g., Hartigan et al. 1995).

Far-ultraviolet (FUV) spectra reveal these diverse processes. Johns-Krull, Valenti, & Linsky (2000) and Calvet et al. (2004) found a correlation between the mass accretion rate and the strength of C IV emission from CTTSs. C IV and other high temperature emission lines are much weaker in non-accreting naked T Tauri stars (NTTSs) than in CTTSs (Johns-Krull et al. 2000; Ardila et al. 2002a). Thus, these lines, which trace gas at 10^5 K, are likely produced by the accretion shocks.

Ardila et al. (2002b) detected wind absorption in the profiles of Mg II lines in *HST*/GHRS spectra of 8 CTTSs, sampling a range of inclinations. They found that the equivalent width of the wind absorption is larger for stars with disks that are observed face-on than for stars with disks observed edge-on, suggesting some collimation in the outflow. Similar P-Cygni profiles of strong lines of O I and C II are typically found in FUV spectra of CTTSs. Fluorescent H₂ emission may trace both warm molecular gas at or near the disk surface (Herczeg et al. 2002) and molecular gas in the surrounding cloud that is shocked as it interacts with stellar outflows (Walter et al. 2003; Saucedo et al. 2003).

In this paper, we present *HST*/STIS and *FUSE* spectra, covering the 912–1700 Å spectral range, of RU Lupi, one of the most heavily veiled¹ CTTSs at optical wavelengths. The spectrum is rich in strong absorption features and in emission lines produced by hot and cool gas. In §2, we describe the properties of RU Lupi. In §3, we describe our *HST*/STIS and *FUSE* observations. In §4, we measure the total hydrogen column density in our line of sight to RU Lupi to estimate the extinction. In §5, we measure the mass accretion rate and present hot emission in lines that are likely produced by the accreting gas. In §6, we show wind absorption in line profiles. Finally, in §7, we analyze variability in the FUV emission from RU Lupi. Our conclusions are presented in §8.

2. Properties of RU Lupi

RU Lupi (HD142560) has some of the strongest accretion signatures among CTTSs, including strong optical veiling and an H α equivalent width that has been observed to vary in the range 140–216 Å (e.g. Appenzeller et al. 1983; Lamzin et al. 1996). Stempels & Piskunov (2002) estimate an optical veiling of 10–20 at 4091 Å and 2–5 between 5300–6300 Å. This strong veiling, the result of a high accretion rate, makes it difficult to estimate stellar properties from the uncertain photospheric emission. Stempels & Piskunov (2002) extracted the optical photospheric spectrum of RU Lupi from high-resolution *VLT*/UVES spectra, and measured a radial velocity of -1.9 ± 1 km s⁻¹. They estimate $T_{eff} = 3950$ K and $\log g = 3.9$. Using these numbers, and an approximate mass of $M = 0.8$

¹ratio of accretion flux to photospheric flux, see Joy (1949); Herbig (1962)

$M_{\odot}$ (Lamzin et al. 1996), they calculate $R = 1.66 R_{\odot}$ and $L = 0.49 L_{\odot}$, assuming a distance of 200 pc. The distance is probably closer to 140 pc (de Zeeuw et al. 1999; Bertout et al. 1999). Based on our extinction estimate $A_V = 0.07$ (see §4), $d = 140$ pc, and the temperature and V-band veiling estimated by Stempels & Piskunov (2002), we calculate a photospheric luminosity $L_{phot} = 0.6 L_{\odot}$ and $R = 1.64 R_{\odot}$. Given these characteristics, the Siess et al. (2000) pre-main sequence stellar evolution tracks indicate that RU Lupi has a mass of $0.6 - 0.7 M_{\odot}$ and an age of 2–3 Myr.

Although most previous studies have assumed that RU Lupi is a single star, Gahm, Petrov, & Stempels (2004) used the *VLT* to detect radial velocity changes of 2.5 km s^{-1} in photospheric absorption lines that are consistent with a period of either 3.7 or 5.6 days. They suggest that these changes are not due to spots, and may instead indicate the presence of a close stellar companion. RU Lupi is variable at ultraviolet and optical wavelengths. A photometric period of 3.7 d was also found by Hoffmeister (1958, 1965), but no period was confirmed in long-term monitoring (see Giovannelli et al. 1995). Hutchinson et al. (1989) and Giovannelli et al. (1990) found that RU Lupi is redder when it is fainter, which suggests that the brightening could be due to either an increase in the mass accretion rate, a decrease in circumstellar extinction, or cool starspots.

Stempels & Piskunov (2002) used optical metallic absorption lines to measure a rotational velocity $v \sin i = 9 \pm 1 \text{ km s}^{-1}$. Rotational velocities of CTTs can be at least $v \sin i = 65 \text{ km s}^{-1}$ (Hartmann & Stauffer 1989; Basri & Batalha 1990; Johns-Krull & Valenti 2001). The measured rotational velocity suggests that either the rotational velocity is small or the rotation axis, and presumably the disk, are observed close to face-on. Ardila et al. (2002a) found that the equivalent width of the Mg II wind absorption in NUV spectra of RU Lupi is the largest of the 8 CTTs in their sample, which we would expect if the wind is collimated and the disk is observed face-on. Blueshifted emission in profiles of the optical [O I], Na I, and Ca II H and K lines suggest that the disk obscures wind emission on the far side of the star, supporting the face-on viewing angle to the star (Stempels & Piskunov 2002). Based on the observations described above, we infer that the disk is probably observed close to face-on. If the 3.7 d period is due to rotation and not a close companion, the rotational broadening of absorption lines would imply an inclination of 24° . Spectro-astrometry of RU Lupi shows that H α emission may be streaming SW and NE from the star (Takami et al. 2001). HH55, located $2'$ SW of the star, may be related to outflows from RU Lupi, which would imply that the disk is at least slightly inclined to our line of sight. The temperature and density of the base of the outflows from RU Lupi were also probed using *HST*/STIS observations of emission in Si III] 1892 Å (Gomez de Castro & Verdugo 2001).

The *ROSAT* count rate of RU Lupi was $\sim 9 \times 10^{-3} \text{ cts s}^{-1}$ in a 15.8 ks PSPC observation (0.1–2.4 keV). We convert this to an X-ray flux of $8 \times 10^{-14} \text{ erg cm}^{-2} \text{ s}^{-1}$, or $L_X = 5 \times 10^{-5} L_{\odot}$ using Neuhaüser et al. (1995).

RU Lupi is associated with a molecular cloud, mapped in ^{12}CO ($J = 2-1$) and ^{13}CO ($J = 1-0$) by Gahm et al. (1993) and in ^{12}CO ($J = 2-1$) by Tachihara et al. (2001). Gahm et al. (1993) found no direct evidence of any molecular outflows from RU Lupi, although they note a possible

connection with HH 55. Both studies found that the radial velocity of the molecular gas surrounding RU Lupi is about -0.4 km s^{-1} .

3. OBSERVATIONS

3.1. HST/STIS

We observed RU Lupi with *HST*/STIS¹ as part of *HST* program 8157 (P.I. F. Walter). The spectra consist of five $\sim 2500 \text{ s}$ exposures with the E140M echelle grating, covering 1150–1710 Å, and a 120 s optical exposure with the G430L grating. We used the $0''.2 \times 0''.06$ aperture to minimize contamination by any off-source emission. The FUV observations use a 1024×1024 pixel CsI MAMA detector, and the optical observations use a CCD. The pixel size on the MAMA detector is $0''.036$ in the dispersion direction (which corresponds to about 0.0142 Å or 3.18 km s^{-1} at 1330 Å) and $0''.029$ in the cross-dispersion direction. Details of these observations are listed in Table 1.

The STIS FUV spectra were obtained in time-tag mode. Figure 1 shows that the count rate is similar at the beginning of each orbit but steadily increases within each orbit by about 40%. Figure 2 shows that the width of the point-spread function in the cross-dispersion direction, measured in regions of strong line emission, decreased during these observations. We conclude that the telescope focus improved during each orbit. Thermal variations of *HST* can change the telescope focus. Models of thermal focus variations³ indicate that the focus was poor at the start of each orbit but improved during every orbit. The nominal FWHM of the point-spread function in the FUV-MAMA is about 2.5 pixels. Because the size of the aperture ($0''.2 \times 0''.06$) in the dispersion direction is smaller than the line-spread function, the count rate increases as the focus improves.

Several other observations obtained using the E140M grating and the $0''.2 \times 0''.06$ aperture on STIS, including those of V471 Tau, a white dwarf with a K2V companion (O’Brien, Bond, & Sion 2001), and several CTTSs also suffer from this problem. On the other hand, observations of the two subluminescent B stars PG1144+615 and PG1219+534 record a constant count rate during each orbit. No other STIS observations were taken in time-tag mode with the $0''.2 \times 0''.06$ aperture. This effect may also be seen data taken with small apertures that are not in time-tag mode.

V471 Tau was observed at three different epochs, twice with the $0''.2 \times 0''.06$ aperture and once with the $0''.2 \times 0''.2$ aperture (see Table 2, O’Brien et al. 2001; Walter 2004). The white dwarf dominates FUV emission and has a 555 s period (Jensen et al. 1986), during which its FUV flux varies by a few percent (Walter 2004). V471 Tau is also a bright FUV source, making it excellent to use for calibration. Figure 3 shows the total flux from 1250–1600 Å in 300 s intervals of four

²See Leitherer et al. (2001) for a description of STIS.

³See <http://www-int.stsci.edu/instruments/observatory/focus/focus2.html#breathing> for a description of modeling the HST focal length variations.

orbits from the observations of V471 Tau. We find one exposure taken with the small aperture, STIS observation number 24906, during which the detected flux is constant, at 4.01×10^{-10} erg cm $^{-2}$ s $^{-1}$. Certain exposures, such as observation numbers 97314 and 97316, show flux increases at a rate about half that found in the observations of RU Lupi. These observations have a peak flux of 3.92×10^{-10} erg cm $^{-2}$ s $^{-1}$. The observations taken with the large aperture have a flux 1250–1600 Å of 4.46×10^{-10} erg cm $^{-2}$ s $^{-1}$.

We measure the flux and cross-dispersion point-spread function in each 300 s interval for the observations of RU Lupi and V471 Tau. The point-spread function depends on wavelength, so we measure the FWHM of emission in the cross-dispersion direction at wavelengths with strong emission lines (O I, C II, Si IV, and C IV) in the spectra of RU Lupi, and in the same regions in the spectra of V471 Tau. Figure 4 shows the percent of the total flux between 1230–1650 Å as a function of point-spread function for V471 Tau and RU Lupi. We calibrate the flux of RU Lupi by calculating a best-fit line to the data from V471 Tau, and then minimizing square of the difference between the data from RU Lupi and the best-fit line calculated from the V471 Tau observations.

For RU Lupi, we calculate a total flux of 6.02×10^{-12} erg cm $^{-2}$ s $^{-1}$ between 1230–1650 Å and 8.43×10^{-13} erg cm $^{-2}$ s $^{-1}$ between 1545–1555 Å (a region dominated by C IV emission). We estimate that this flux calibration has an absolute uncertainty of about 15%. Inspection of the V471 Tau observations indicate that the changes in flux are constant with wavelength, such that the relative flux calibration across the wavelength range is uncertain to $< 3\%$ within a single exposure and between different exposures. The consistency of the relative count rate in strong lines during our observation of RU Lupi indicates that the relative flux is uncertain by $< 10\%$.

The Doppler correction in our data was in error by a factor of 1.6, such that the spectrum shifts by 3.8 pixels (12 km s $^{-1}$) during a single orbit. We corrected for this error by subdividing the spectra into 300 s intervals, and subsequently shifting and coadding the spectra. This error is prevalent in STIS echelle spectra in the FUV, but is now corrected in newer versions of the *calstis* pipeline.⁴ The spectral resolution is about $R = 45,000$.

The wavelengths were initially calibrated using short exposures of the on-board Pt/Cr-Ne lamp obtained at the end of each observation of RU Lupi. We determined the absolute wavelength calibration independently by measuring the wavelength of the geocoronal Ly α emission line. We then shift the line to 1215.58 Å, where we expect the line to be at the time of our observation. We find that our wavelength solution corresponds well to that based on the calibration lamp. The absolute and relative wavelength solution is accurate to at least 3 km s $^{-1}$.

⁴see http://www.stsci.edu/hst/stis/calibration/pipe_soft_hist/update215c.html

3.2. FUSE

We obtained a 25 ks spectrum of RU Lupi with the *FUSE* satellite in 21 separate exposures, each covering 912–1187 Å, as part of *FUSE* program A109 (P.I. F. Walter). These observations occurred at night to minimize contamination by airglow emission. The *FUSE* has four co-aligned telescopes. The optical paths are coated with SiC or Al:LiF to maximize efficiency between 900–1000 Å or 1000–1187 Å, respectively. One observation produces eight overlapping spectra, each of which covers about 90 Å. A detailed description of the *FUSE* mission can be found in Moos et al. (2000) and Sahnou et al. (2000). Since we used the LWRS aperture ($30'' \times 30''$), the *FUSE* spectra would include emission extended beyond the point source of RU Lupi, if present.

The exposures were coadded and reduced with the *calFUSE* v2.4 software package (Kruk, Murphy, & Young 2001). The 8 separate spectra were then coadded using *hrs_merge* in *IDL*. No significant differences were found in overlapping wavelength ranges of these segments. Since the LiF channels are more sensitive than the SiC channels, the S/N in the *FUSE* spectrum is higher at $\lambda > 1000$ Å. The data were binned by 4 pixels, which is smaller than the approximate instrument resolution of 9 pixels, or $R \approx 15,000$. The flux scale is accurate to about $\pm 5\%$, including a small degradation in some of the channels over time.

We calibrate wavelengths in our *FUSE* spectrum by comparing the measured wavelengths of H₂ emission lines (see §5.2) in the *FUSE* spectrum to those in the STIS spectrum. The H₂ lines in the STIS spectrum can be characterized by a bright narrow component and a fainter and broader blueshifted component. When convolved to the lower spectral resolution of *FUSE*, the STIS H₂ lines appear Gaussian with a velocity shift of -15.5 km s^{-1} relative to the photospheric radial velocity of RU Lupi. We offset the pipeline-calibrated wavelengths by $+5 \text{ km s}^{-1}$, so that the average H₂ emission line in the *FUSE* spectrum has a velocity shift of -15.5 km s^{-1} . This velocity shift also places the peak of the N I 1135 Å emission line at the radial velocity of the star, which is similar to the velocity shift of emission lines from neutral atoms in the STIS spectrum (see §5.2). The shift of 5 km s^{-1} is less than the 15 km s^{-1} accuracy of the standard *calFUSE* pipeline wavelength calibration (Kruk, Murphy, & Young 2001). We estimate an uncertainty of about 5 km s^{-1} in our wavelength calibration at $\lambda > 1100$ Å, and a larger uncertainty at $\lambda < 1100$ Å.

The STIS spectrum overlaps the *FUSE* spectrum between 1160–1185 Å, although the sensitivity of STIS is poor at these short wavelengths. Figure 5 shows that the C III line at 1175 Å is much stronger in the STIS spectrum than in the *FUSE* spectrum, so fluxes from the two spectra are not directly comparable.

4. THE ISM IN OUR LINE OF SIGHT TO RU LUPI

4.1. Interstellar Absorption Lines

Narrow absorption lines, produced in the interstellar medium (ISM), are detected against several stellar emission lines (see Table 3 and Figure 6). In §5.2, we find that most low-ionization emission lines have Gaussian profiles centered about the radial velocity of RU Lupi. We therefore assume that the intrinsic emission profile is Gaussian to measure the velocity and equivalent width of the absorption.

Interstellar absorption is centered between -8 and -19 km s $^{-1}$ in the heliocentric frame, depending on the line. RU Lupi is located at $l = 338^\circ, b = +11^\circ 89'$, where models of the local interstellar medium produced by Redfield & Linsky (2000) predict that very little absorption by the LIC is expected and some absorption by the G cloud is possible. These models also predict that any absorption in the local interstellar cloud and the G cloud should be centered at -23 km s $^{-1}$ and -26 km s $^{-1}$, respectively. The Lupus molecular cloud has a velocity of about -0.4 km s $^{-1}$ in our line of sight toward RU Lupi (Gahm et al. 1993; Tachihara et al. 2001). Tachihara et al. (2001) estimate that the H $_2$ column density of the Lupus cloud in the direction of RU Lupi is $\log N(\text{H}_2) = 21.0$, which is much higher than the column density $19.0 < \log N(\text{H}_2) < 19.5$ in our line of sight to RU Lupi, suggesting that the star is located in front of most of the cloud. The data show absorption components redshifted relative these clouds, indicating that other interstellar clouds are located along the RU Lupi line of sight.

Ardila et al. (2002b) found that the equivalent widths of interstellar Mg II lines towards CTTSs are larger than typical values measured through the ISM toward nearby sources. We also detect large equivalent widths in many narrow absorption lines in the FUV, including a detection of interstellar absorption in the Si III 1206.5 Å line. Interstellar Si III absorption has only been detected in one other local sightline, towards ϵ CMa (Gry et al. 1995; Wood et al. 2002b). The large Mg II equivalent widths and detection of Si III absorption may be produced by warm or shocked gas in the Lupus molecular cloud. Any additional absorption components due to the Lupus molecular cloud, a remnant envelope, or a slow wind should be centered within a few km s $^{-1}$ of the interstellar absorption.

Ly α is detected in emission only longward of 1217.5 Å, indicating that the sum of interstellar and circumstellar H I absorption is optically thick shortward of 1217.5 Å. We do not detect any blueshifted emission from Ly α because of strong H I absorption in the wind. ISM absorption should dominate absorption on the red side of the line. We model the observed Ly α profile with a single Gaussian emission profile, centered at the radial velocity of the star, and a Voigt absorption profile. The height of the Gaussian is determined from the best fit to the red side of the Ly α line for a given $N(\text{H I})$. We assume that the absorption is centered at -15 km s $^{-1}$ in the heliocentric frame, roughly the velocity at which we detect interstellar absorption in lines such as O I and C II (Table 3). We assume a Doppler broadening parameter $b \sim 11$ km s $^{-1}$, assuming a temperature of 7000

K, typical of the local ISM (Redfield & Linsky 2000). At the high column densities applicable to the RU Lupi sightline, the column density is insensitive to b . Figure 7 compares the profiles that best fit the observed emission for a range of neutral hydrogen column density $N(\text{H I})$. We find that $\log N(\text{H I}) = 20.0 \pm 0.15$ (cm^{-2} units), where the error includes an estimate of the uncertainty in the central velocity of the ISM absorption.

The low S/N in the continuum prevents us from detecting most of the H_2 absorption lines that should occur in the *FUSE* bandpass. However, the 6-0 P(3) line at $1031.19 \text{ \AA}$ is detected because it occurs against strong O VI emission (Fig. 8). This H_2 line is shifted by about $-23 \pm 5 \text{ km s}^{-1}$ relative to the stellar radial velocity. The O VI $1037 \text{ \AA}$ line is weak and narrow because most of the line is absorbed by H_2 and C II (see Fig. 8). The detection of the one H_2 line and only weak O VI $1037 \text{ \AA}$ emission imply that $19.0 < \log N(\text{H}_2) < 19.5$ for temperatures $100 \text{ K} < T(\text{H}_2) < 300 \text{ K}$, which is typical of interstellar molecular hydrogen (Rachford et al. 2002). We therefore measure a total hydrogen column density of $\log N(\text{H}) = \log[N(\text{H I}) + 2 \times N(\text{H}_2)] = 20.1 \pm 0.2$. Stempels & Piskunov (2002) measured optically thin absorption in the Na I line at $5890 \text{ \AA}$, and found $\log N(\text{Na I}) = 11.0$. They used interstellar abundances from Welty & Hobbs (2001) to estimate $\log N(\text{H}) \sim 19.8$, which is approximately what we measure.

4.2. Dust Extinction

Strong optical veiling due to accretion prevents a reliable estimate of the extinction using photospheric emission from RU Lupi. Instead, we use the interstellar absorption discussed above to constrain the extinction. Assuming a typical interstellar gas/dust ratio and dust grains described by $R_V = 3.09$ (Cardelli, Clayton, & Mathis 1989), we convert the hydrogen column density of $\log N(\text{H}) = 20.1 \pm 0.2$ to an extinction $A_V \sim 0.07 \text{ mag}$ using the relation described by Bohlin et al. (1978). This low extinction implies that RU Lupi is near the front edge of the Lupus molecular cloud. We adopt $A_V = 0.07 \text{ mag}$ for the remainder of this paper. We confirm a low extinction to RU Lupi using two other probes in our data.

The NUV and blue continuum of RU Lupi is produced by accretion (e.g. Hartigan et al. 1991). We use the shape of the accretion continuum to constrain the extinction by modelling the continuum as emission from an isothermal slab, described in §5.1. Model continua that best fit the observed UV-optical ($1650 \text{ \AA} < \lambda < 5700 \text{ \AA}$) continuum are calculated for each assumed value of the extinction. Figure 9 shows the estimated accretion continuum, calculated independently for three assumed extinctions compared with the extinction-corrected NUV spectrum of RU Lupi. When we use $A_V > 0.4 \text{ mag}$, our models of the accretion continuum underestimate the flux at $2175 \text{ \AA}$ because the extinction bump at $2175 \text{ \AA}$ increases the assumed intrinsic flux at that wavelength.

If instead most of the dust grains in our line of sight to RU Lupi are in the Lupus molecular cloud, then the dust may differ from typical interstellar dust. The grain size should not differ significantly from grains typical of the local ISM. Although sightlines through the densest regions

of molecular clouds can have $R_V = 5.5$, Kenyon, Dobrzycka, & Hartmann (1994) and Whittet et al. (2001) find that the sightlines through the Taurus and Ophiucus molecular clouds with $A_V < 3.0$ do not vary significantly from the average interstellar value of $R_V = 3.09$ (Cardelli et al. 1989). However, the extinction law towards the peculiar B star HD29647 ($A_V = 3.6$ mag), which lies behind Taurus, does not show the strong 2175 Å bump (Whittet et al. 2004) that is typically associated with small carbonaceous dust grains (e.g. Draine & Malhotra 1993). The grains in the line of sight to HD29647 have an $R_V = 3.63$, which suggests that they are large relative to typical interstellar grains, described by $R_V = 3.09$, (Cardelli et al. 1989). Calvet et al. (2004) applied the HD29647 extinction law to analyze ultraviolet spectra of intermediate-mass T Tauri stars in Taurus. If the extinction law of HD29647 applies to low-extinction targets such as RU Lupi, then the shape of the accretion continuum is not a good probe of extinction because of the absence of the 2175 Å bump.

We also place an upper limit on the extinction by comparing the relative strength of fluorescent H₂ emission lines (see §5.2), which occur between 1100–1650 Å, to the same emission lines from TW Hya. Extinction is negligible to TW Hya, and both stars are observed nearly face-on, so any geometrical effects on the H₂ fluxes should be similar. In §5.2, we identify 90 unblended H₂ lines in the STIS spectrum of RU Lupi from 15 different upper levels. In the STIS spectrum of TW Hya, Herczeg et al. (2002) detected 146 H₂ lines from 19 different upper levels. By comparing the relative flux in lines from the same upper level for the two stars, we place an upper limit of $A_V < 0.6$ mag toward RU Lupi for either a typical interstellar extinction law or the interstellar extinction law of HD29647.

Our estimate of a low extinction towards RU Lupi agrees with most previous estimates. Gahm et al. (1979) estimated $0.3 < A_V < 1.0$ mag using the *IUE* spectrum of RU Lupi. Giovannelli et al. (1995) estimated an extinction of $A_V = 0.3$ mag based on the 2200 Å reddening bump in *IUE* spectra, and their result agrees with the extinction $A_V = 0.4$ mag estimated by Brooks et al. (2001) and Lamzin et al. (1996). Our extinction estimate is much lower the extinction of $A_V = 1.28$ mag calculated by Hughes et al. (1994) based on optical colors, which are contaminated by emission due to accretion.

The molecular fraction of gas towards RU Lupi is between 17–40%. Most sightlines with a molecular fraction > 0.01 have $A_V > 2.4$ mag (Savage et al. 1977; Rachford et al. 2002). However, this relationship may not apply to environments around young stars. A high molecular fraction and low extinction were also found towards the Herbig Ae star HD104237 (Grady et al. 2004).

5. ACCRETION

5.1. Mass Accretion Rate

We use the accretion luminosity, estimated from the UV excess and the Balmer jump at 3700 Å, to approximate the mass accretion rate of RU Lupi (Valenti et al. 1993; Gullbring et al. 1998). Our *HST* observations consist of nearly simultaneous optical and FUV spectra. We obtain NUV spectra of RU Lupi using a combination of archival observations from *HST*/STIS, *HST*/GHRS, and *IUE* (Table 4). The NUV spectra are normalized by factors listed in Table 4 to match the optical and FUV continua flux, ignoring regions of strong emission lines. We scale the optical STIS spectrum of V836 Tau, a K7 WTTS, to use as a template for the photospheric emission from RU Lupi. We estimate that $A_V = 1.52$ mag towards V836 Tau, the average extinction estimate from Kenyon & Hartmann (1995) and Whittet et al. (2001).

The methodology used to calculate the mass accretion rate is described by Valenti, Basri, & Johns (1993) and Herczeg et al. (2004). We model the spectrum of the accretion shock by approximating the emission from an isothermal, pure-hydrogen slab (see Fig. 10), which fit the data well. Other studies have also calculated the accretion continuum by modelling the accretion shock (Calvet & Gullbring 1998). The emission spectrum that best fits the observed spectra is produced by a slab with temperature $T = 9600$ K, electron density $n_e = 3 \times 10^{14} \text{ cm}^{-3}$, slab surface area parameter (fraction of stellar surface area covered by accretion) $\delta = 0.017$, thickness $l = 2 \times 10^7$ cm, and a scaling factor for the template photospheric spectrum of 0.39. The slab has an optical depth of 0.64 at 5000 Å, which is less than the average optical depth in slab models of blue emission from other CTTSs (Basri & Bertout 1989; Hartigan et al. 1991). This scaling factor corresponds to an optical veiling in the continuum of a factor of 4–8 between 4000–4300 Å, and about 2 at longer wavelengths. We calculate an accretion luminosity $L_{slab} = 0.35 L_\odot$. We then estimate a mass accretion rate of

$$\dot{M}_{acc} = \frac{1.25 R_* L_{slab}}{GM_*} = (5 \pm 2) \times 10^{-8} \frac{d^2}{140^2} \frac{R}{1.65 R_\odot} \frac{0.8 M_\odot}{M} M_\odot \text{yr}^{-1} \quad (1)$$

This mass accretion rate is large, but significantly smaller than previous rates that were estimated assuming $A_V = 1.3$ (e.g., Hughes et al. 1994; Ardila et al. 2002a). Although mass accretion rates can be as high as $10^{-6} M_\odot \text{yr}^{-1}$ for CTTSs, the mass accretion rate of $5 \times 10^{-8} M_\odot \text{yr}^{-1}$ is high for CTTSs with ages of 2–3 Myr (Hartmann et al. 1998).

The model accretion continuum accounts for the observed optical and NUV continuum. A slight FUV excess may be detected, although the model accretion continuum in the FUV is sensitive to uncertainties in the temperature of the accreting gas. Figure 11 shows that the FUV continuum from RU Lupi, if present, is much weaker than that detected towards TW Hya and several other CTTSs (Herczeg et al. 2004; Bergin et al. 2004). The FUV continuum from CTTSs is most likely produced by H₂ fluorescence generated by highly energetic electrons colliding with and electronically exciting H₂ molecules, and may be related to an inner dust hole in the disk (Liu et al. 2002; Bergin et al. 2004). A bump at 1600 Å, diagnostic of this H₂ emission, is weakly present.

5.2. FUV Emission Lines

The total observed emission between 1170–1700 Å from RU Lupi is 7.16×10^{-12} erg cm $^{-2}$ s $^{-1}$, most of which occurs in emission lines. The flux in hot FUV emission lines from CTTSs increases with the mass accretion rate (Johns-Krull et al. 2000; Calvet et al. 2004). Many Fe II and fluorescent H $_2$ lines are pumped by other emission lines, such as Ly α , so their strength should also scale with the accretion rate. Johns-Krull et al. (2000) found that H $_2$ flux scales with the C IV flux. Other lines, such as the O I 1305 Å triplet and the C II 1335 Å doublet, may be produced both in accreting gas and in the wind. Thus, we can analyze certain emission lines to probe the accretion flow.

For emission lines that are not corrupted by wind absorption, we fit a single or multiple Gaussians to the emission line profile to measure the line center, FWHM, and flux in the line. Tables 5 and 6 list parameters for identified atomic or unidentified emission lines detected in the STIS and *FUSE* spectra, respectively. For lines that have large wind absorption components, we list only the continuum-subtracted flux detected in the line. Fluxes are not directly comparable between the STIS spectrum and the *FUSE* spectrum owing to temporal variability in the emission (see Fig. 5).

Strong resonance lines, such as the C IV doublet at 1549 Å, are commonly detected in emission from late-type stars and are also strong in the FUV spectrum of RU Lupi. We identify weak emission lines using the atomic spectral line database of Kurucz & Bell (1995). The identifications marked by a ‘?’ in Tables 5–6 are uncertain. Since many lines, including fluorescent H $_2$ emission, Fe II, and Cr II, are pumped by Ly α , our identification technique relies on searching for lines that could be pumped by Ly α or other strong emission lines. We list the pumped lines and the pumping transition in Table 7. Many of these lines were identified by Johansson & Jordan (1984), Carpenter et al. (1988), and Hartman & Johansson (2000). Other pumped lines include Cl I 1351.6 Å, which is pumped by C II 1336 Å, and S I 1472 Å, which is pumped by O I 1302 Å and S I 1295 Å. The O I triplet at 1304 Å can be pumped by Ly β 1026 Å (Carpenter et al. 1988). Although airglow Ly β prevents us from detecting the center of Ly β in our *FUSE* spectrum, we do not detect Ly β emission at 100 km s $^{-1}$. Nevertheless, the O I lines are very strong. Moreover, other lines such as C II that require similar excitation are also very strong. We speculate that the O I emission in the RU Lupi spectrum is most likely produced by collisional excitation or recombination in the hot gas rather than by fluorescence.

Many narrow (FWHM $\approx$ 17.5 km s $^{-1}$) emission lines detected in the STIS spectrum are identified as H $_2$ lines (see Table 8) based on the H $_2$ transition database calculated by Abgrall et al. (1993) and the linelist from *HST*/STIS observations of TW Hya (Herczeg et al. 2002). Most of these lines are Lyman-band (B-X) H $_2$ lines pumped by Ly α . We also detect one Werner-band (C-X) H $_2$ line at 1208.8 Å that is pumped by O VI emission. These H $_2$ lines are found in UV spectra of CTTSs (Brown et al. 1981; Valenti et al. 2000; Ardila et al. 2002a), red giants (Ayres, Brown, & Harper 2003a), and the wind of Mira B (Wood et al. 2002a). We coadded the strongest H $_2$ lines in the STIS

spectrum to analyze the line profiles. Figure 12 shows that the coadded H_2 lines are well fit by a strong Gaussian, centered at -12 km s^{-1} , and a weaker Gaussian, centered at -30 km s^{-1} , relative to the radial velocity of RU Lupi. We also detect additional flux in these lines from $v = -120$ to -70 km s^{-1} . The flux ratio of these two Gaussians is 1.3 and the ratio of the widths is 0.33. To measure fluxes, we fit each individual H_2 line with two Gaussians, with the velocity separation, relative width, and relative flux fixed to the numbers obtained from the fit to the coadded profile above. We also detect unresolved Lyman and Werner-band H_2 lines with *FUSE*. We fit a single Gaussian profile to these lines, and list their fluxes in Table 9. If present, any extended H_2 emission would be detected by *FUSE* because it has a large ($30'' \times 30''$) aperture. Figure 13 compares the coadded spectral line profiles of the H_2 lines in the *STIS* and *FUSE* spectra, and the geocoronal $\text{Ly}\beta$ line in the *FUSE* spectrum. The *FUSE* H_2 lines are not significantly broader than the *STIS* H_2 lines, convolved to the lower resolution of *FUSE*. If the H_2 emission were extended beyond a point source in the dispersion direction, we would expect the spectral line profile would be much wider. This comparison therefore suggests that the H_2 emission in the *FUSE* spectrum is not significantly extended beyond a point source. The H_2 lines will be discussed in detail in a future paper.

Some cooler emission lines typically found in stellar chromospheres, such as the O I 1304 Å triplet and the C II 1335 Å doublet, are partially or completely absorbed by wind features listed in Table 10 (see also Fig. 16). Figure 14 shows that emission in these cool lines are centered at the radial velocity of RU Lupi.

Emission in C IV, Si IV, N V, and possibly O VI is most likely produced by 10^5 K gas heated by the accretion shock. Figure 15 shows that these lines are strong and broad. The *STIS* lines have centroids near the radial velocity of the star. The *FUSE* lines of O VI 1032 Å and C III 977 Å lines are blueshifted by ~ 190 and $\sim 100 \text{ km s}^{-1}$, respectively. Low S/N prevents any detection of wind in the C III 977 Å line, but if present we may be underestimating the blueshift. Absorption in H_2 and other lines may mask some redshifted emission in the O VI and C III lines. The velocity shift of these lines may be time dependent. Alternatively, the emission in C III and O VI resonance lines during the *FUSE* observation might be produced by shocked emission due to the outflow, as hot gas can be seen in some HH objects (Raymond et al. 1997). The lack of C III 1175 Å emission suggests that any emission produced by accreting gas is faint during this observation. *FUSE* would then only detect this spatially extended blueshifted emission. The *STIS* observations would most likely not be sensitive to this extended emission because of its small aperture. The time dependence in the strength of C III emission is discussed in §7.

The C IV 1548 Å line shows broad blueshifted emission, extending to -300 km s^{-1} , that is not present in the other emission lines. In contrast, the Si IV 1394 Å line has stronger redshifted emission than the Si IV 1403 Å line. The difference in behavior between the C IV doublet and the Si IV doublet is puzzling. The N V 1243 Å line is absorbed by N I in the wind (Fig 16), while interstellar H_2 absorption corrupts the O VI line profiles (Fig. 8). Anomalously weak N V 1243 Å emission was also detected in the FUV spectrum of T Tau (Walter et al. 2003).

Optical depth effects likely produce these complex emission features. The mass accretion rate is large, so we infer that the accretion column may be optically thick. The C IV 1548 Å and 1550 Å lines from TW Hya have a similar shape to each other (Herczeg et al. 2002), possibly because the mass accretion rate of TW Hya is 25 times smaller than that of RU Lupi, so the accretion shock may be less optically thick. We also note that, unlike TW Hya, Si II, Si III, and Si IV lines from RU Lupi are all bright. Understanding these profiles will require detailed modelling of the accreting gas.

Most of the observed Ly α emission is absorbed by H I in the stellar outflows, in the ISM, and possibly the accretion shock itself. In Herczeg et al. (2004), we found that the intrinsic Ly α emission could be reconstructed from the H₂ lines, and may contribute 85% of the total FUV emission from CTTs. The strength of the H₂ and Fe II lines that are pumped by Ly α imply that the Ly α emission from RU Lupi is similarly strong. We will estimate the Ly α emission from RU Lupi in a future paper.

Coronal Fe lines have been detected in UV spectra of the Sun (Sandlin, Brueckner, & Tousey 1977) and cool stars (Ayres et al. 2003b; Redfield et al. 2003). Lamzin et al. (2001) predict that emission in the [Fe XII] 1349.382 Å and Fe XI 1467.420 Å lines may be produced at the accretion shock and could be detected in UV spectra of CTTs. Based on models of the accretion shock by Lamzin et al. (2001), the lines should be centered at $v_{shock}/4$. The FWHM of these emission lines should be dominated by thermal broadening. We do not detect any coronal Fe lines in the RU Lupi spectra, and place unreddened upper limits of $\sim 2 \times 10^{-15}$ erg cm⁻² s⁻¹ that are about an order of magnitude larger than the line fluxes predicted by Lamzin et al. (2001).

6. WIND ABSORPTION

Outflows from RU Lupi are detected by the strong blueshifted absorption seen in P-Cygni profiles and against the continuum (Fig. 6). The absorption is typically seen from low energy levels of neutral or singly-ionized species.

The cool emission lines that are not corrupted by wind absorption are symmetric about the radial velocity of the star (see §5.2 and Figure 14). We therefore estimate the intrinsic profile of lines with wind absorption by reflecting the red emission about the radial velocity of the star. We then measure the equivalent width and central and maximum velocities of the absorption feature (see Table 10). Lines such as O I, C II, and Fe II have P-Cygni profiles, with absorption ranging from about -70 to -270 km s⁻¹. For certain lines of Fe II and N I, the wind absorption is detected against the continuum. The wind absorption feature seen in Si III is narrower than the wind absorption features detected in other lines from the ground state, which may be due either to a small column density of Si III or the absence of a velocity gradient for the hot gas in our line of sight. The latter possibility could occur if most of the Si III occurs close to the star. Most of the wind absorption is detected in transitions with low-energy lower levels. However, we also detect

wind absorption in the N I lines at 1242, 1492, and 1494 Å from lower levels excited to 2 eV (Fig. 16).

In certain cases, the wind absorption is strong enough that many nearby lines are barely detected. Figure 17 shows the O I triplet at 1304 Å and the C II doublet at 1335 Å. The wind in the C II 1335.7 Å line obscures emission in C II 1334.5 Å. Similarly, the O I 1305 Å line is weak due to wind in that line and a wind in O I 1306 Å.

7. VARIABILITY

Figure 5 shows a striking decrease in flux in the C III 1175 Å multiplet between the STIS and the *FUSE* observations. In the *FUSE* spectrum, most of the emission detected near 1175 Å is H₂ and not C III. The O VI 1035 Å doublet and C III 977 Å emission are weak and blueshifted. If the drop in flux seen in C III 1175 Å is related to a decrease in the mass accretion rate, then we may also not be detecting any emission produced by the accretion shock in the VI 1035 Å doublet or the C III 977 Å line. Instead, with its large aperture, *FUSE* may have detected emission in these lines that is produced by shocks in the outflow. The fluorescent H₂ emission in the *FUSE* spectrum, which is highly sensitive to the Lyα emission, is comparable to the flux we expect based on the STIS spectrum. In §5.2, we found that the H₂ emission in the *FUSE* bandpass originates predominantly on-source. Because the H₂ lines serve as a proxy for the strength of Lyα emission, we infer that the Lyα emission has a similar flux in the two observations. This result is surprising because we expect that both C III and Lyα emission are produced by accreting gas. We speculate that a smaller mass loss rate significantly reduces the optical depth of the C III line, while the Lyα line remains optically thick. The FUV continuum where the STIS and *FUSE* spectra overlap is marginally detected at $\sim 2 \times 10^{-15}$ erg cm⁻² s⁻¹ Å⁻¹. This emission, which is likely produced by H₂ (Bergin et al. 2004), did not change by more than $\sim 50\%$.

We also compare the observed STIS spectrum with spectra obtained using the G160M grating on *HST*/GHRS, which cover narrow (35 Å) spectral regions centered on the C IV and Si IV lines (Ardila et al. 2002a). In the GHRS spectra, the FUV emission lines have similar shapes but are four times weaker than the fluxes detected with STIS (Fig. 18). The continuum in the GHRS observation also appears weaker than that detected in the STIS observation. However, the H₂ emission line fluxes are similar in the STIS and the GHRS spectra, which suggests that the Lyα emission did not change significantly between the two observations. The GHRS observation could also include off-source H₂ emission, as is detected towards T Tau (Walter et al. 2003). Our STIS observations must have fortuitously occurred when the accretion rate of RU Lupi was higher than usual.

We find no significant variability during the STIS spectrum. Although this analysis is complicated because the count rate depends on the changing focus of *HST* (see §3.1), we place a limit that the flux of RU Lupi varied by no more than 10% in intervals longer than 100 s and 4% in inter-

vals longer than 300 s. Some minor differences appear in the profiles of hot lines during the STIS observations. The S/N of the *FUSE* observations prevented any significant analysis of variability over short timescales.

8. CONCLUSIONS

We present and analyze FUV spectra of RU Lupi, a 2–3 Myr old CTTS with a high mass accretion rate. Based on these spectra, we find the following results:

1. The *HST* telescope focus improved during each orbit of the RU Lupi observation, which increased the detected count rate because the FWHM of the line-spread function was larger than the aperture size in the dispersion direction. We account for this instrumental problem by using observations of V471 Tau. This problem appears in several other time-tag STIS spectra that used a small aperture.

2. We estimate a total hydrogen column density $\log N(\text{H}) = 20.1 \pm 0.2$ in our line of sight to RU Lupi using Ly α and H₂ absorption, and convert the column density to an extinction of $A_V \sim 0.07$ mag.

3. The mass accretion rate of RU Lupi is about $(5 \pm 2) \times 10^{-8} M_{\odot} \text{ yr}^{-1}$.

4. Emission is detected in 174 lines, 103 of which are Lyman and Werner-band H₂ lines pumped by Ly α or O VI. Some Fe II and Cr II lines are also pumped by Ly α . The hot lines of C IV, Si IV, and N V are bright and broad, but differences in the line profiles are puzzling. In particular, the C IV 1548 Å line has strong redshifted emission that is not seen in the C IV 1550 Å line, but the Si IV 1394 Å line has strong blueshifted emission relative to the Si IV 1403 Å line. Emission in the O VI 1035 Å doublet and C III 977 Å is blueshifted, and at the time of our *FUSE* observation may be dominated by emission produced by shocks in the outflow. The observed Ly α emission only accounts for about 1% of the observed FUV emission, likely because of strong absorption by H I in the ISM and stellar wind.

5. Wind absorption is detected in 30 different lines, and generally ranges from -70 to -270 km s⁻¹. Atoms with low first-ionization potentials are ionized in the wind, resulting in absorption in lines of Si II, Si III, Fe II, Al II, and C II. Wind absorption is also detected in lines of N I and O I. Three N I wind absorption lines originate from levels excited 2.38 eV above the ground state.

6. The FUV emission from RU Lupi varies significantly over months-long timescales. Our STIS observation of RU Lupi occurred during a period of strong emission, possibly due to a high mass accretion rate. The C III emission is much weaker in the *FUSE* spectrum than in the STIS spectrum, although the H₂ emission, which is a proxy for Ly α emission, does not vary significantly between these two observations.

9. ACKNOWLEDGEMENTS

This research was funded in part by STScI grant GO-08157.01-97A to SUNY Stony Brook, by STScI grant GO-08157.02-97B to the University of Colorado, NASA contract NNG04GJ31G from Johns Hopkins University to the University of Colorado, and by the Swedish National Space Board. This paper is based on observations made with the NASA/ESA Hubble Space Telescope, obtained at the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS5-26555, and on observations obtained by the NASA-CNES-CSA FUSE mission. FUSE is operated for NASA by the Johns Hopkins University under NASA contract NAS5-32985. These observations are associated with *HST* program 8157 and *FUSE* program A109. We thank the anonymous referee for insightful comments that improved the paper. We also thank Paul Goudfrooij for suggestions and discussion concerning the instrumental problems with *HST*/STIS. GH thanks Brian Wood and Seth Redfield for discussion related to the calibration of the STIS spectra.

REFERENCES

- Abgrall H., Roueff, E., Launay, F., Roncin, J. Y., & Subtil, J. L. 1993, A&AS, 101, 273
- Appenzeller, I., Krautter J., & Jankovics, I. 1983, ApJ, 308, 836
- Ardila, D.R., Basri, G., Walter, F.M., Valenti, J.A., Johns-Krull, C.M. 2002a, ApJ, 566, 1100
- Ardila, D.R., Basri, G., Walter, F.M., Valenti, J.A., Johns-Krull, C.M. 2002b, ApJ, 567, 1013
- Ayres, T.R., Brown, A., Harper, G.M. 2003a, ApJ, 598, 610
- Ayres, T.R., Brown, A., Harper, G.M., Osten, R.A., Linksy, J.L., Wood, B.E., & Redfield, S. 2003b, ApJ, 583, 963
- Basri, G., & Batalha, C. 1990, ApJ, 363 654
- Basri, G., & Bertout, C. 1989, AJ, 341, 340
- Bergin, E. et al. 2004, ApJL, 614, 143
- Bertout, C., Robichon, N., & Arenou, F. 1999, A&A, 352, 574
- Bohlin, R.C., Savage, B.D., & Drake, J.F. 1978, ApJ, 224, 132
- Brooks, D.H., Costa, V.M., Lago, M.T.V.T., & Lanzafame, A.C. 2001, MNRAS, 327, 177
- Brown, A. Jordan, C., Millar, T. J., Gondhalekar, P., & Wilson, R. 1981, Nature, 290, 34
- Calvet, N. & Gullbring, E. 1998, ApJ, 509, 802

- Calvet, N., Muzerolle, J., Briceno, C., Hernandez, J., Hartmann, L., Saucedo, J.L., & Gordon, K.D. 2004, *ApJ*, accepted
- Cardelli, J. A., Clayton, G. C., & Mathis, J. S. 1989, *ApJ*, 345, 245
- Carpenter, K.G., Pesce, J.E., Stencel, R.E., Brown, A., Johansson, S., & Wing, R.F. 1988, *ApJS*, 68, 345
- Draine, B.T., & Malhotra, S. 1993, *ApJ*, 414, 632
- Gahm, G.F., Fredga, K., Liseau, R., & Dravins, D. 1979, *A&AL*, 73, 4
- Gahm, G.F., Johansson, L.E.B., & Liseau, R. 1993, *A&A*, 274, 415
- Gahm, G.F., Petrov, P.P., & Stempels, H.C. 2004, *Proc. Cool Stars, Stellar Systems and the Sun* 13, ESA SP, in press.
- Giovannelli, F., Rossi, C., Castaldo, D., Covino, E., & Errico, L. 1990, *Ap&SS*, 169, 55
- Giovannelli, F. et al. 1995, *A&AS*, 114, 341
- Gomez de Castro, A.I., & Verdugo, E. 2001, *ApJ*, 548, 976
- Grady, C. et al. 2004, *ApJ*, 608, 809
- Gry, C., Lemonon, L., Vidal-Madjar, A., Lemoine, M., & Ferlet, R. 1995, *A&A*, 302, 497
- Gullbring, E., Hartmann, L., Briceno, C., & Calvet, N. 1998, *ApJ*, 492, 323
- Hartigan, P., Edwards, S., & Ghandour, L. 1995, *ApJ*, 452, 736
- Hartigan, P., Kenyon, S.J., Hartmann, L., Strom, S.E., Edwards, S., Welty, A.D., & Stauffer, J. 1991, *ApJ*, 382, 617
- Hartman, H., & Johansson, S. 2000, *A&A*, 359, 627
- Hartmann, L., & Stauffer, J. R. 1989, *AJ*, 97, 873
- Hartmann, L., Calvet, N., Gullbring, E., & D'Alessio, P. 1998, *ApJ*, 495, 385
- Herbig, G.H. 1962, *AdA&A*, 1, 47
- Herczeg, G. J., Linsky, J. L., Valenti, J.A., Johns-Krull, C.M. 2002, *ApJ*, 572, 310
- Herczeg, G.J., Wood, B.E., Linsky, J.L., Valenti, J.A., Johns-Krull, C.M. 2004, *ApJ*, 607, 369
- Hoffmeister, C. 1958, *Veröff Stenwarte Sonneberg*, 3, 333
- Hoffmeister, C. 1965, *Veröff Stenwarte Sonneberg*, 6, 91

- Hughes, J., Hartigan, P., Krautter, J., & Kelemen, J. 1994, *AJ*, 108, 1071
- Hutchinson, M.G., Evans, A., Davies, J.K., & Bode, M.F. 1989, *MNRAS*, 237, 683
- Jensen, K.A., Swank, J.H., Petre, R., Guinan, E.F., Sion, E.M., & Shipman, H.L. 1986, *ApJL*, 309, 27
- Johansson, S., & Jordan, C. 1984, *MNRAS*, 210, 239
- Johns-Krull, C. M., Valenti, J. A., & Linsky, J. L. 2000, *ApJ*, 539, 815
- Johns-Krull, C. M. & Valenti, J. A. 2001, *ApJ*, 561, 1060
- Joy, A.H. 1949, *ApJ*, 110, 424
- Kenyon, S.J., Dobrzycka, D., & Hartmann, L. 1994, *AJ*, 108, 1872
- Kenyon, S.J., & Hartmann, L. 1995, *ApJS*, 101, 117
- Kruk, J., Murphy, E. & Young, P. 2001, *The Caluse Pipeline Reference Guide*
- Kurucz, R. L. & Bell, B. 1995, CD-ROM No. 23. (Cambridge: Smithsonian Astrophysical Observatory)
- Lamzin, S.A., Stempels, H.C., & Piskunov, N.E. 2001, *A&A*, 369, 965
- Lamzin, S.A., Bisnovatyi-Koga, G.S., Errico, L., Giovannelli, F., Katysheva, N.A., Rossi, C., & Vittone, A.A. 1996, *A&A*, 306, 877
- Leitherer, C., et al. 2001, *STIS Instrument Handbook, Version 5.1* (Baltimore: STScI)
- Lindler, D. 1999, *CALSTIS Reference Guide* (Greenbelt: NASA/LASP)
- Liu, X., Shemansky, D. E., Abgrall, H., Roueff, E., Dziczek, D., Hansen, D. L., & Ajello, J. M. 2002, *ApJS*, 138, 229
- Moos, W.H., et al. 2000 *ApJ*, 538, L1
- Neuhaüser, R., Sterzik, M.F, Schmitt, J.H.M., Wichmann, R., & Krautter, J. 1995, *A&A*, 297, 391
- O’Brien, M. S., Bond, H. E., & Sion, E. M. 2001, *ApJ*, 563, 971
- Rachford, B. L., Snow, T. P., Tumlinson, J., Shull, J. M., Blair, W. P., Ferlet, R., Friedman, S. D., Gry, C., et al. 2002, *ApJ*, 577, 221
- Raymond, J.C., Blair, W.P., Long, K.S. 1997, *ApJ*, 489, 314
- Redfield, S. & Linsky, J.L. 2000, *ApJ*, 534, 825

- Redfield, S., Ayres, T.R., Linsky, J.L., Ake, T.B., Dupree, A.K., Robinson, R.D., Young, P.R. 2003, ApJ, 585, 993
- Sahnou, D.J., et al. 2000, ApJ, 538, L7
- Sandlin, G.D., Brueckner, G.E., & Tousey, R. 1977, ApJ, 214, 898
- Saucedo, J., Calvet, N., Hartmann, L., & Raymond, J. 2003, ApJ, 591, 275
- Savage, B.D., Bohlin, R.C., Drake, J.F., & Budich, W. 1977, ApJ, 216, 291
- Siess L., Dufour E., & Forestini M. 2000, A&A, 358, 593
- Stempels H.C., & Piskunov, N. 2002, A&A, 391, 595
- Tachihara, K., Toyoda, S., Onishi, T., Mizuno, A., Fukui, Y., & Neuhäuser, R. 2001, PASJ, 53, 1081
- Takami, M., Bailey, J., Gledhill, T.N., Chrysostomou, A., & Hough, J.H. 2001, MNRAS, 323, 177
- Valenti, J.A., Basri, G., & Johns, C.M. 1993, ApJ, 106, 2024
- Valenti, J. A., Johns-Krull, C. M., & Linsky, J. L. 2000, ApJS, 129, 399
- Valenti, J.A., Fallon, A.A., & Johns-Krull, C.M. 2003, ApJS, 147, 305
- Walter, F.M., et al. 2003, AJ, 126, 3076
- Walter, F.M. 2004, AN, 325, 241
- Welty, D.E., & Hobbs, L.M. 2001, ApJS, 133, 345
- Whittet, D. C. B., Gerakines, P. A., Hough, J.H., & Shenoy, S. S. 2001, ApJ, 547, 872
- Whittet, D.C.B., Shenoy, S.S., Clayton, G.C., & Gordon, K.D. 2004, ApJ, 602, 291
- Wilkinson, E., Harper, G.M., Brown, A., Herczeg, G.J. 2002, AJ, 124, 1077
- Wood, B. E., Karovska, M. & Raymond, J. C. 2002a, ApJ, 575, 1057
- Wood, B. E., Redfield, S., Linsky, J.L., & Sahu, M.S. 2002b, ApJ, 581, 1168
- de Zeeuw, P.T., Hoogerwerf, R., de Bruijne, J.H.J, Brown, A.G.A., & Blaauw, A. 1999, AJ, 117, 354




Fig. 1.— In our STIS observations of RU Lupi, the count rate within each orbit increased with time, but then returned to a low level at the start of the next orbit. This increase of flux results because the FWHM of the line-spread function is wider than the aperture in the dispersion direction but narrows during each orbit, due to thermal changes in the telescope.




Fig. 2.— The FWHM of C IV and Si IV emission in the cross-dispersion direction, measured in 300 s intervals, improves during each orbit because of thermal focus variations.




Fig. 3.— The flux detected between 1250–1600 Å in four of 16 orbits of STIS observations of V471 Tau. A gradual increase of detected flux with time is present in two of the orbits shown here, and is similar to the increase seen in our spectra of RU Lupi.




Fig. 4.— The flux between 1230–1650 Å from RU Lupi (diamonds) and V471 Tau (asterisks) depends on the telescope focus. The fraction of the total flux is calculated by comparing the measured flux in a 300 s interval to the estimated total flux, for the wavelength region between 1230–1650 Å. The image size in the dispersion direction is plotted here as the average of the FWHM of O I (1292–1313 Å), C II (1333–1337 Å), Si IV (1392–1405 Å), and C IV (1547–1552 Å). The dotted line is the line that best fits the data from the calibration source V471 Tau. We then estimate the total flux from RU Lupi by fitting the RU Lupi data to this line using the least squares minimization method. The point-spread function is broader than the small aperture, so that better focus improves the count rate on the detector increases.




Fig. 5.— C III emission was strong during the HST/STIS observation (dotted line), but was much weaker during the *FUSE* observation (solid line), which occurred about a year later. This striking difference likely resulted because of a drop in the mass accretion rate between the two observations. The wavelengths of fluorescent H₂ emission lines are indicated by the dashed vertical lines.




Fig. 6.— P-Cygni profiles are detected for many lines, four of which are shown here. The wind absorptions in most lines range from $\sim -70 - 270 \text{ km s}^{-1}$, although the absorption in the Si III line is much narrower than wind absorption from ground state transitions of other species because the line is less optically thick. Narrow absorption near line center (except for the excited O I 1306 Å line) is produced by the interstellar medium and possibly also by material local to RU Lupi. Interstellar Si III absorption is not detected locally (Wood et al. 2002b). The dashed vertical line marks the average velocity of interstellar absorption detected towards RU Lupi.




Fig. 7.— The observed red wing of the $\text{Ly}\alpha$ line (solid line) is modeled as a Gaussian profile, centered at the heliocentric velocity of $\text{Ly}\alpha$, and a Voigt absorption profile for interstellar H I, centered at the interstellar velocity measured from other absorption lines (see Table 5). No $\text{Ly}\alpha$ emission is detected on the blue side of line center. We show here the fit for three different assumed column densities of H I. We determine that $\log N(\text{H I}) \approx 20.0 \pm 0.15$.




Fig. 8.— The *FUSE* spectrum of RU Lupi (solid line) is compared with the transmission percentage (dashed line) through gas with $T = 300$ K and $\log N(\text{H}_2)=19.0$. Both lines in the O VI doublet are blueshifted from line center (dotted vertical lines). The O VI 1032 Å profile shows strong emission, with absorption in the H₂ line at 1031.19 Å. The O VI 1037 Å line is detected, but most of the line is absorbed by C II (dashed vertical lines) and H₂ in the line of sight to the star.




Fig. 9.— We fit models (dashed lines) of the accretion continuum to the observed UV-optical spectrum of RU Lupi, assuming a range of extinctions. When we assume $A_V > 0.4$ mag, we cannot reconcile the increased emission at 2175 Å with the emission at shorter wavelengths. Low extinction is also consistent with the observed $N(\text{H I})$ column density in our line of sight to the star (see §4.1).




Fig. 10.— We model the observed accretion continuum (solid line) with emission from a pure hydrogen slab (dashed line). The accretion continuum is obtained from data described in Table 3, assuming $A_V = 0.07$. To extract the accretion continuum shown here, we subtracted the scaled spectrum of the WTTS V819 Tau, a template for the photospheric emission of RU Lupi, from the observed spectrum of RU Lupi.




Fig. 11.— Comparison of the observed UV emission from RU Lupi (bottom solid line) and TW Hya (top solid line, multiplied by 10) with spectra calculated from modelling the accretion continuum (dashed lines). The FUV continuum in excess of emission from the accretion column is much weaker from RU Lupi than from TW Hya.




Fig. 12.— The spectral profile of coadded H₂ lines from the STIS spectrum (solid line) shows that the lines are blueshifted and asymmetric. We fit the profile with two Gaussian profiles (dotted). The summed profiles are shown by the dashed line, with parameters described in §5.2. Excess blueshifted emission at $v = -120$ to -70 km s⁻¹ is also detected. These lines will be discussed in detail in a future paper.




Fig. 13.— We compare the spectral profile of coadded H₂ lines observed with *FUSE* (solid line) with geocoronal Lyβ emission (dotted line) and coadded H₂ lines from the STIS spectrum (dashed line), convolved to the spectral resolution of *FUSE*. The geocoronal Lyβ emission fills the aperture and in this plot is shifted in velocity to the center of the H₂ lines. The coadded *FUSE* H₂ lines have a similar FWHM as the coadded STIS H₂ lines, which suggests that the H₂ emission detected with *FUSE* is dominated by on-source emission. Any emission extended in the dispersion direction would broaden the line profile.




Fig. 14.— The weighted mean of the radial velocity of cool (neutral, singly, and doubly ionized) lines is -1.4 km s^{-1} (dotted line) relative to the radial velocity of RU Lupi. Lines that are contaminated by wind emission and lines that are only tentatively identified (marked by a ? in Table 5) are not used in this analysis.




Fig. 15.— Emission lines produced in $\sim 10^5$ K gas are broad and strong in spectra of CTTSs. These lines from RU Lupi show complicated features. The C IV 1548 Å line shows a strong blue bump that is not detected in any of the other lines, while the Si IV 1394 Å line shows excess redshifted emission relative to the other lines. N V 1243 Å emission is absorbed by N I in the wind (see Fig. 16). The O VI and C III lines detected in the *FUSE* spectrum appear blueshifted and may be produced by shocks in the outflow. Because the *FUSE* observation occurred about one year after the STIS observation and the FUV emission from RU Lupi is variable, the O VI and C III lines shown above should not be directly compared with the C IV, Si IV, or N V lines.




Fig. 16.— Wind absorption is detected in N I lines from excited lower levels, with $E=2.38$ eV. The N I absorption near 1243 Å attenuates the N V 1243 Å emission line.




Fig. 17.— Wind absorption in the O I and C II lines. The wind absorbs a significant amount of the emission line and also emission from adjacent lines. O I 1305 $\text{\AA}$ is weak because of wind absorption due to O I 1306 $\text{\AA}$. Similarly, C II 1334.5 $\text{\AA}$ is barely detected because of absorption in C II 1335.7, and two H₂ lines are weak because of absorption in C II 1334.5. Line wavelengths at the radial velocity of RU Lupi are indicated by the vertical dotted lines. The narrow absorption feature near line center is produced by the ISM.




Fig. 18.— The STIS spectrum of RU Lupi has a stronger FUV continuum and stronger emission lines than the previously observed GHR spectrum (Ardila et al. 2002a).

Table 1: OBSERVATIONS OF RU LUPI

Date	Instrument	Exposure (s)	Bandpass (Å)	Aperture	Spectral Res. ($\lambda/\Delta\lambda$)
7/12/00 ^a	<i>HST</i> /STIS E140M	12530	1150–1700	0′.2 × 0′.06	45000
7/12/00 ^a	<i>HST</i> /STIS G430L	120	2900–5700	52′ × 0′.1	800
9/28/01 ^b	<i>FUSE</i>	24517	900–1187	30′ × 30′	15000
^a JD 2451738		^b JD 2452180			

Table 2: *HST*/STIS OBSERVATIONS OF V471 TAU

Dates	Orbits	Exposure	Aperture	F_{1400}^a	Program	Obs. Numbers ^b	P.I.
3/13–3/23/98 ^c	6	14350	0′.2 × 0′.06	4.01	7735	24906–24919	F. Walter
8/24/00–8/27/00 ^d	4	7848	0′.2 × 0′.2	4.46	8305	73220–73229	H. Bond
1/25/02 ^e	4	10336	0′.2 × 0′.06	4.01	9283	97310–97316	F. Walter

^aPeak flux from 1250–1600 Å, in 10^{-10} erg cm⁻² s⁻¹

^bSTIS observation numbers

^cJD 2450885–2450896

^dJD 2451781–2451784

^eJD 2452299

Table 3: Properties of Interstellar Absorption Lines^a

ID	λ_{obs} (Å)	v (km s ⁻¹)	W_λ (Å)
Si III	1206.447	-11.2	0.089 ± 0.007
O I	1302.090	-16.2	0.186 ± 0.008
C II	1335.662	-8.3	0.098 ± 0.005
Si II	1526.618	-15.1	0.172 ± 0.009
Fe II	1608.367	-13.7	0.107 ± 0.010
Al II	1670.672	-18.7	0.173 ± 0.008

^aMeasured by assuming Gaussian emission lines.

Table 4: SPECTRA OF THE ACCRETION CONTINUUM

Bandpass	Instrument	Scaling Factor ^a	S/N ^b	Date
1185–1700	<i>HST</i> /STIS E140M	1.0	20	07/12/00
1700–1839	<i>IUE</i> SWP-LO	0.75	60	many ^c
1839–1930	<i>HST</i> /STIS G230M	1.1	35	08/24/92
1930–2302	<i>IUE</i> SWP-LO	0.75	14	many ^c
2302–2350	<i>HST</i> /GHRS G270M	1.1	40	08/24/92
2350–2775	<i>IUE</i> SWP-LO	0.75	70	many ^c
2775–2822	<i>HST</i> /GHRS G270M	1.1	120	08/24/92
2900–5700	<i>HST</i> /STIS G430L	1.0	100	07/12/00

^aScaling factor to normalize flux to the *HST* fluxes.

^bApproximate S/N in a 5 Å bin.

^cMultiple *IUE* spectra were coadded by (Valenti, Fallon, & Johns-Krull 2003) and are available at <http://archive.stsci.edu/prepds/iuepms/datalist.html>.

Table 5: Properties of emission lines in the STIS spectrum^a

ID	λ_{obs} Å	v km s ⁻¹	FWHM km s ⁻¹	Flux ^b	ID	λ_{obs} Å	v km s ⁻¹	FWHM km s ⁻¹	Flux ^b
C III	1175.164	-	327	57.1	Fe II	1360.112	-12	128	9.5
C III	1176.563	-	103	14.4	Fe II	1366.344	-9	197	12.0
Si II	1190.662	64	134	3.5	Fe II	1368.653	-32	102	3.2
Si II	1194.462	-8	75	3.7	Fe II?	1379.474	3	210	8.0
Si III (1) ^d	1206.149	-85	105	24.8	Si IV (1) ^d	1393.860	25	310	240
Si III (2) ^d	1206.495	1	362	214	Si IV (2) ^d	1393.861	25	158	150
H I ^c	1217.500	-	-	75	Si IV (1) ^d	1402.636	-27	113	78
N V (1) ^d	1238.151	-160	368	28.0	Si IV (2) ^d	1402.704	-12	275	225
N V (2) ^d	1238.828	4	135	20.3	P I?	1411.783	12	111	5.7
?	1252.154	-	81	3.0	Fe II?	1414.920	10	172	4.5
Si II ^c	1265.000	-	-	11.3	Cr II	1428.624	-7	54	2.0
C I?	1280.956	28	77	1.5	Fe II	1464.918	-11	118	5.3
Fe II	1288.641	-17	228	5.8	S I	1471.847	5	115	13.7
Si III	1294.451	-20	127	10.1	C I?	1481.645	-22	211	7.5
S I	1295.610	-8	100	22.3	S I?	1483.426	21	213	12.2
S I	1296.056	-25	260	46.9	Fe II	1506.572	10	127	8.7
Si III	1298.992	13	203	22.8	Si II ^c	1526.000	-	-	14.2
O I ^c	1302.000	-	-	91	Si II ^c	1533.000	-	-	20
O I ^c	1304.762	-20	47	33.3	C IV (1) ^d	1547.900	-54	202	173
O I ^c	1306.000	-	-	137	C IV (2) ^d	1548.092	-16	435	268
Si II ^c	1309.261	-1	257	59	C IV (3) ^d	1548.287	21	35	31.5
P II	1309.870	1	106	18.8	C IV (1) ^d	1550.807	9	229	192
Si II ^e	1310.747	12	232	16.0	C IV (2) ^d	1550.839	15	65	40
Fe II ^f	1312.630	34	238	5.6	Fe II ^c	1558.928	-28	161	7.2
S I	1323.355	-34	141	4.2	Fe II ^c	1563.543	-45	91	6.1
S I	1326.572	-14	92	1.5	Fe II ^c	1566.486	35	211	13.3
C I	1329.518	-11	189	5.2	Fe II	1631.005	-17	144	12.9
Fe II?	1330.985	9	146	4.8	He II (1) ^d	1640.455	-1	256	35
C II ^c	1334.000	-	-	6.3	He II (2) ^d	1640.499	7	70	28
C II ^c	1335.000	-	-	112	He II (3) ^d	1641.284	150	124	12.7
Cr II	1347.136	24	83	2.9	O III]	1666.228	16	237	47
Cl I	1351.649	0	135	27.2	Al II ^c	1670.000	-	-	15
O I	1355.615	6	157	35.5	Cr II?	1697.215	-34	184	15.5
C I?	1357.240	25	93	2.0	Fe II ^c	1702.000	-	-	23.6
O I	1358.438	-14	88	5.5					

^aLines fit with single Gaussian, except where noted.

^cWind absorption corrupts the emission line profile.

^ePossibly blended with Si I, Si II emission.

^fPossibly blended with S I

^b10⁻¹⁵ erg cm⁻² s⁻¹

^dFit with multiple Gaussians labeled (1), (2), or (3)

Table 6: Atomic and unidentified Emission Lines in the *FUSE* spectrum^a

ID	λ_{obs} (Å)	v (km s ⁻¹)	FWHM (km s ⁻¹)	Flux ^b
C III (1) ^{c,d}	976.154	-263	106	3.7
C III (2) ^{c,d}	976.679	-102	95	10.5
?	997.313	-	36	0.8
O VI	1031.251	-190	359	30.4
O VI ^e	1036.861	-215	52	2.8
?	1048.151	-	67	2.3
?	1074.385	-	42	1.3
N I ^c	1134.284	-33	128	5.4
N I ^c	1134.974	0	64	3.3
N I?	1168.514	-4	79	1.5

^aLines fit with single Gaussian.

^b10⁻¹⁵ erg cm⁻² s⁻¹

^cWind absorption corrupts emission profile.

^dFit with multiple Gaussians.

^eCorrupted by H₂ absorption.

Table 7: Pumped atomic lines in the STIS spectrum

ID	λ_{obs} (Å)	ID _{pump}	λ_{pump} (Å)
Fe II	1288.641	H I	1213.738
S I	1295.610	O I	1302.336
S I	1296.056	O I	1302.862
Si II	1309.261	O I	1304.370
P II	1309.870	O I	1301.874, 1304.675
P II	1310.747	O I	1305.497
Fe II	1312.630	H I	1215.873
Fe II?	1330.985	O I	1304.436
Cr II	1347.149	H I	1215.765
Cl I	1351.651	C II	1335.726
Fe II	1360.117	H I	1215.503
Fe II	1366.345	H I	1214.735
Fe II	1368.649	H I	1214.735
Fe II?	1379.474	Si IV?	1403.255
P I?	1411.783	H I	1216.608
Fe II?	1414.920	C III	1175.147
Cr II	1428.624	H I	1217.387
Fe II	1464.918	H I	1213.738
S I	1471.840	O I	1302.336
Fe II	1506.544	C II	1335.409
Cr II?	1697.215	H I	1304.381

Table 8: Identification of H₂ lines in the STIS spectrum

ID	λ_{obs} (Å)	Flux ^a	ID	λ_{obs} (Å)	Flux ^a	ID	λ_{obs} (Å)	Flux ^a
1-2 R(3)	1202.396	3.1	1-4 R(6)	1327.506	1.7	1-7 R(3)	1489.497	10.6
1-5 R(3) ^b	1208.787	1.8	1-4 P(5)	1329.047	-1.0	4-8 P(8)	1494.917	1.2
0-2 P(3)	1225.469	2.1	0-4 P(2)	1338.512	13.0	1-7 R(6)	1500.364	8.6
2-2 R(11)	1237.511	1.2	0-4 P(3)	1342.192	12.9	1-7 P(5)	1504.679	18.5
1-2 P(8)	1237.827	4.5	4-6 R(3)	1359.031	1.4	4-9 R(3)	1513.489	1.7
4-4 R(3)	1253.625	1.5	4-6 R(6)	1371.055	0.7	3-7 R(15)	1513.916	1.5
3-1 P(17)	1254.074	1.7	4-4 R(17)	1372.014	0.9	blend ^c	1516.122	4.6
1-3 R(3)	1257.780	6.4	4-6 P(5)	1372.671	1.3	0-7 P(2)	1521.529	1.8
3-2 R(15)	1265.121	0.7	2-4 P(13)	1379.913	1.1	1-7 P(8)	1524.586	8.8
4-4 R(6)	1266.515	0.6	1-5 P(5)	1387.299	3.4	0-7 P(3)	1525.092	1.5
4-4 P(5) ^c	1266.823	1.7	0-5 P(1)	1396.157	0.9	2-8 R(11)	1555.810	3.2
3-1 P(18)	1268.806	0.5	0-5 P(2)	1398.883	10.8	1-8 R(6)	1556.796	7.5
3-4 P(1)	1270.527	0.6	2-5 R(11)	1399.151	1.8	1-8 P(5)	1562.317	6.9
1-3 R(6)	1270.967	3.1	0-5 P(4)	1407.154	3.2	4-10 P(5)	1572.548	1.6
2-2 P(13)	1271.128	2.4	blend ^d	1408.807	1.8	1-8 P(8)	1580.573	4.8
1-3 P(5)	1271.869	5.1	3-5 R(15)	1418.169	1.1	0-8 P(2)	1582.350	3.1
4-2 R(17)	1273.968	1.7	1-6 R(3)	1430.945	9.6	2-8 P(13)	1588.709	6.7
0-3 R(0)	1274.495	6.4	2-5 P(13)	1434.439	2.2	3-10 P(1)	1591.263	1.3
0-3 R(1)	1274.855	4.5	1-6 R(6)	1442.794	5.8	3-9 R(15)	1593.191	3.9
0-3 R(2)	1276.222	1.0	1-6 P(5)	1446.051	14.2	2-9 R(11)	1602.176	4.9
3-2 R(16)	1277.958	0.6	2-6 R(11)	1453.024	3.4	4-11 R(3)	1602.543	3.0
0-3 P(2)	1279.407	5.9	0-6 R(0)	1454.761	3.7	1-9 R(3)	1603.177	2.5
0-3 P(3)	1283.051	6.1	0-6 R(1)	1454.894	5.1	4-11 R(6)	1606.202	2.1
0-3 P(4)	1287.632	0.4	0-6 R(2)	1455.847	1.0	1-9 R(6)	1610.884	2.5
4-4 P(8)	1287.886	0.6	0-6 P(1)	1457.348	1.1	4-11 P(5)	1613.645	4.8
2-3 R(11)	1290.842	0.8	0-6 P(2)	1460.095	7.6	1-9 P(5)	1617.817	2.3
1-3 P(8)	1293.814	4.4	0-6 P(3)	1463.712	11.6	3-9 P(17)	1622.084	1.5
1-4 R(3)	1314.570	2.6	4-6 R(17)	1465.096	2.1	4-11 P(8)	1622.821	3.8
3-2 P(18)	1320.133	0.7	1-6 P(8)	1467.014	6.2	2-9 P(13)	1632.458	3.1
3-5 P(1)	1324.966	0.8	4-8 P(5)	1476.996	1.0	3-11 P(1)	1636.275	3.9

^a10⁻¹⁵ erg cm⁻² s⁻¹

^bWerner-band (C-X), pumped by O VI at 1031.862 Å

^cAlso includes emission from 4-1 P(19)

^dBlend of 1-5 P(8) and 3-4 P(17)

^eBlend of 0-1 R(0) and 0-1 R(1)

Table 9: Lyman and Werner-band H₂ lines in the *FUSE* spectrum

ID ^a	λ_{obs} (Å)	v (km s ⁻¹)	FWHM (km s ⁻¹)	Flux ^b
C-X 2-2 P(5)	1058.255	-22	31	1.3
B-X 1-0 P(5)	1109.273	-8	36	0.4
C-X 1-3 Q(3)	1119.017	-14	71	0.7
C-X 0-2 Q(10)+ C-X 1-3 P(5)	1127.096	-	61	1.5
B-X 1-1 R(3)	1148.632	-16	30	1.4
C-X 1-3 P(11)	1154.850	-14	36	1.5
B-X 1-1 P(5)+ B-X 1-1 R(6) ^c	1161.774	-	60	3.4
B-X 0-1 P(2)	1166.176	-18	34	0.9
B-X 0-1 P(3)	1169.687	-15	38	0.8
C-X 0-3 Q(10) + C-X 1-4 P(5)	1171.916	-	47	3.4
blend ^d	1174.396	-	122	4.9
C-X 2-6 R(3)	1174.786	-19	25	1.6
C-X 1-3 Q(16)	1177.208	-14	21	0.6

^aB-X refers to Lyman-band, C-X refers to Werner-band

^b10⁻¹⁵ erg cm⁻² s⁻¹

^cB-X 0-1 R(0) and B-X 0-1 R(1) may also contribute

^cC-X 1-5 Q(7), C-X 1-4 R(9), C-X 2-5 R(0), C-X 2-5 R(1), C-X 2-5 R(2)

Table 10: Properties of Wind Absorption Lines^a

ID	λ_{obs} (Å)	v_c^b (km s ⁻¹)	v_{max}^c (km s ⁻¹)	E''^d (cm ⁻¹)	W_λ (Å)
Si II	1189.75	-170	-290	0	1.20 ± 0.07
N I	1199.83	-	-	0	1.06 ± 0.11^e
Si III	1206.75	-180	-220	0	0.20 ± 0.02
N I	1242.74	-110	-190	19224	0.86 ± 0.04
Si II	1265	-150	-	287	1.15 ± 0.15^f
O I	1301.43	-170	-290	0	0.94 ± 0.01
Si II	1303.57	-180	-260	0	0.61 ± 0.10^g
O I	1304.30	-130	-220	158	0.72 ± 0.10^g
O I	1305.30	-170	-280	227	0.88 ± 0.01
Si II	1308.55	-170	-270	287	0.78 ± 0.05
C II	1334.5	-140	-230	0	0.98 ± 0.25
C II	1334.93	-170	-300	63	1.08 ± 0.01
Cl I	1346.35	-200	-320	0	0.38 ± 0.10
S I	1424.60	-120	-210	0	0.27 ± 0.04
N I	1492.02	-120	-210	19224	0.77 ± 0.05
N I	1493.75	-190	-310	19224	0.56 ± 0.03
Si II	1525.84	-170	-280	0	0.96 ± 0.05
Si II	1532.55	-170	-280	287	0.97 ± 0.05
Fe II	1558.18	-170	-270	1872	0.60 ± 0.09
Fe II	1562.93	-160	-260	2430	0.50 ± 0.05
Fe II	1565.95	-70	-140	1872	0.40 ± 0.10
Fe II	1574.23	-150	-200	1872	0.23 ± 0.05
Fe II	1579 59	-190	-290	2430	0.53 ± 0.04
Fe II	1607.55	-170	-270	0	0.66 ± 0.08
Fe II	1612.79	-190	-270	1872	0.55 ± 0.09
Fe II	1620.73	-170	-230	385	0.34 ± 0.04
Fe II	1624.64	-160	-260	2430	0.47 ± 0.06
Fe II	1628.15	-180	-250	667	0.28 ± 0.07
Al II	1669.85	-170	-270	0	1.02 ± 0.06
Fe II	1701.22	-140	-200	1872	0.53 ± 0.05

^aMeasured assuming an emission line symmetric about the radial velocity of the star or as absorption against the continuum.

^bCentral velocity of absorption.

^cMaximum velocity of absorption.

^dEnergy of lower level.

^eMultiplet.

^fUncertain blue edge

^gBlend